

Are you a first time visitor? Be sure to register.

Search



→ HOME

→ ABOUT NURSA

→ REGISTRATION

→ DATASETS

→ REAGENTS

E-JOURNAL

▪ Editorial Information and  
Instructions for Authors

→ MEETINGS

→ FEEDBACK

→ SEARCH

→ LINKS

→ HELP

# NRS | NUCLEAR RECEPTOR SIGNALING

The Open Access Journal of the Nuclear Receptor Signaling Atlas

NURSA Home : Nuclear Receptor Signaling

&lt;&lt; Previous Article | Next Article &gt;&gt;

Volume 1 | 2003

Review

Cite this Article: Nuclear Receptor Signaling (2003) 1, e008.

## RNAi technology and its use in studying the function of nuclear receptors and coregulators

Hai-Jun Zhou, Sophia Y. Tsai and Ming-Jer Tsai

Department of Molecular and Cellular Biology, Baylor College of Medicine, One  
Baylor Plaza, Houston, Texas 77030

Received: July 16, 2003; Accepted: August 26, 2003; Published: September 10, 2003

Copyright © 2003, Zhou et al. This is an open-access article distributed under the terms of the  
Creative Commons Non-Commercial Attribution License, which permits unrestricted non-commercial  
use distribution and reproduction in any medium, provided the original work is properly cited.  
Article DOI: 10.1621/nrs.01008

### Abstract

### Article Navigation

Until just a few years ago, RNA interference (RNAi) technology was restricted to the research fields of plants, *C. elegans* or *Drosophila*. The discovery of gene silencing by in vitro synthesized double-stranded RNA (dsRNA) in mammalian cells has made the use of RNAi possible in nearly the entire life science kingdom. DNA vectors delivering small interfering RNA (siRNA) directed by polymerase III or polymerase II promoters to persistently inhibit target genes expression have extended this technology to study in vivo function of these genes. Recently, RNAi has been used as a powerful tool in the functional analysis of nuclear receptors and their coregulators. This short review will cover studies in this area.

### Abbreviations

RNAi: RNA interference; shRNA: Short hairpin RNA; siRNA: Small interfering RNA

### Introduction

### Article Navigation

RNA interference (RNAi) was first discovered in the nematode worm *Caenorhabditis elegans* as a biological response to double-stranded RNA (dsRNA), which resulted in potent sequence-specific gene silencing [Fire et al., 1998]. RNAi is an evolutionarily conserved process involving a multi-step event, which generates small interfering

RNAs (siRNAs) of 21- to 23-nucleotide (nt) *in vivo* by endogenous RNase III enzyme-Dicer. The resulting siRNAs mediate destruction of their complementary mRNA. The biology and mechanisms of RNAi have been reviewed recently in detail [Fire et al., 1998; Hannon, 2002]. The breakthrough of RNAi study is the discovery that dsRNA can selectively suppress gene expression in cultured mammalian cells through RNAi [Caplen et al., 2001; Elbashir et al., 2001a]. The short (<30 nt) synthetic interfering RNA duplexes are successfully used to induce sequence-specific gene silencing yet evade the host interferon response which usually is activated by dsRNA longer than 30nt [Baglioni and Nilsen, 1983; Williams, 1997]. Based on the biochemical analysis of siRNA in *Drosophila* [Elbashir et al., 2001b; Elbashir et al., 2001c; Zamore et al., 2000], the structure of the *in vitro* synthesized siRNA is found to be important to achieve effective gene inhibition [Elbashir et al., 2001a]. siRNAs with 3' overhangs of two uridines have been found to be more efficient in cultured mammalian cells [Elbashir et al., 2001c]. At the moment, there is no criterion to predict the ideal target sequence of an siRNA. Although many of the siRNAs reported to date are designed to target coding sequences, especially the amino terminus 100-200 bases away from AUG [Sui et al., 2002], successful gene silencing has been reported for siRNA by targeting various sequences, including the 3' untranslated region [McManus et al., 2002]. Therefore, the target sequences published in the literatures will definitely provide a candidate pool for scientists who are interested in the RNAi research.

A major drawback of using *in vitro* synthesized siRNAs is its transient nature because mammalian cells lack the mechanism to amplify siRNA-mediated silencing observed in *C. elegans* and *Drosophila* [Hannon, 2002]. Gene expression was only suppressed for no more than one week. In the beginning of 2002, DNA vectors expressing siRNAs directed by RNA polymerase III promoters were made by several groups [Brummelkamp et al., 2002; Paddison et al., 2002; Sui et al., 2002]. These expression vectors mediate the production of siRNAs from transcripts containing a stem and loop structure-short hairpin RNA (shRNA), which will lead to the continued expression of siRNAs in the cells with a persistent and specific knock-down of the target genes. This improved expression system paves the way for long-term loss-of-function studies.

However, the Polymerase III promoter is active in all tissues and cannot be used to generate the tissue-specific knock-down. Just recently, a new vector expressing long ds-RNA from the Polymerase II promoter has been developed to knock-down the target gene [Shinagawa and Ishii, 2003]. The ds-RNA transcribed from this vector lacking 5'-cap structure and 3'-poly(A) tail, which facilitate its exportation to cytoplasm, can silence target gene without producing the interferon response. Transgenic mice embryos expressing long ds-RNA for the transcriptional co-repressor Ski from this vector exhibited similar phenotypes to those of Ski-knock-out embryos. Therefore, with this polymerase II expression vector, one can efficiently knock down the expression of any gene in animal in a tissue-specific manner without the host interferon response.

#### RNAi for nuclear receptors and coregulators

##### Article Navigation

RNA interference with synthesized dsRNA or vector-based siRNAs has endowed research with a whole set of tools that facilitate genetic studies of many fields, including the recent application of RNAi approaches in functional analysis of nuclear receptors and their co-regulators. Since last year, RNAi has been introduced to silence the expression of nuclear receptors and their co-regulators in transient transfection assays (see Table 1 for details). FXR [Plass et al., 2002] and AR [Wright et al., 2003] are the only published examples of nuclear receptor superfamily which have been successfully inhibited with synthetic siRNA. For the nuclear receptor co-regulators, RNAi has been more widely used. By introduction of

[illegible]

Abbreviations: FXR, Farnesoid X receptor; AR, Androgen Receptor; SRC-1/2/3, Steroid Receptor Coactivator-1/2/3; NR, Nuclear Receptor Co-repressor; SMRT, Silencing mediator for Retinoid and thyroid hormone receptor; H3AC3, Histone acetylase 3; TBL1, Transducin  $\beta$ -Like protein 1; R1, Transducin  $\beta$ -Like protein 1-Related protein; ER $\alpha$ , Estrogen Related Receptor  $\alpha$ ; ARA55, AR-associated protein 55; A70, AR-associated protein 70. \* Ho-Geun Yoon and Jiemin Wong, personal communication. \*\* Haijun You unpublished results

## Article Navigation

Distinct from the antisense oligodeoxynucleotides (asODN) and dominant negative forms, which mediate RNaseH cleavage and functional interference respectively, RNAi leads to target gene mRNA degradation through the RNAase III machinery resulting in the knock-down of protein expression, thereby affecting the functions of nuclear receptors and coregulators. Although RNAi technology has not been applied in any *in vivo* system of nuclear receptors or coregulators, RNAi induced effects in *in vitro* studies are comparable or better than those produced by antisense oligodeoxynucleotides [Cavarretta et al., 2002] or dominant negative form [Rahman et al., 2003a; Rahman et al., 2003b]. In these studies, asODN to SRC-1 reduced SRC-1 protein level by 78% at best. In contrast, RNAi to SRC-1 is able to reduce 80%-90% of its protein level. In addition, ARA55 and ARA70 knocking-down with DNA vector-based RNAi, reduced the expression of AR target

gene, PSA, to a similar extent as that reduced by ARA55 and ARA70 dominant negative mutants. For the reporter assay, RNAi knocking-down of ARA55 and ARA70 exerted a greater inhibition of MMTV promoter activity than those produced by dominant negative forms of ARA55 and ARA70. In general, the effectiveness in most cases (listed in Table 1) is dramatic and has greatly helped the functional analysis of these proteins.

## Conclusion

### Article Navigation

There is no golden principle to predict the length of sustaining time of RNAi inhibition, because each protein has its own turnover time. Normally, the transient effect of siRNA inhibition lasts less than one week. Multiple transfections may be needed to extend its effect to a longer time. Otherwise, vector-based RNAi either by polymerase II or III is required to archive permanent knock-down of target gene expression. The length of targeting sequences range from 19nt to 21 nt, which guarantee the sequence specificity, without generating interferon response and mimic the product length of Dicer. Usually, the effective sequence is chosen from several candidates; therefore, the targeting sequences listed here will be very helpful for the functional analysis of nuclear receptors and co-regulators. Although RNAi technology has not been used as widely as it can be in the nuclear receptor field, we believe the importance and effectiveness of this technology will be increasingly used for functional analysis of nuclear receptor and coregulator function for the years to come.

## References

### Article Navigation

Baglioni, C. and Nilsen, T. W. (1983) Mechanisms of antiviral action of interferon. *Interferon* **5**, 23-42. [PubMed](#) | [Back to Text](#)

Brummelkamp, T. R., Bernards, R. and Agami, R. (2002) A system for stable expression of short interfering RNAs in mammalian cells. *Science* **296**, 550-3. [PubMed](#) | [Back to Text](#)

Caplen, N. J., Parrish, S., Imani, F., Fire, A. and Morgan, R. A. (2001) Specific inhibition of gene expression by small double-stranded RNAs in invertebrate and vertebrate systems. *Proc Natl Acad Sci U S A* **98**, 9742-7. [PubMed](#) | [Back to Text](#)

Cavarretta, I. T., Mukopadhyay, R., Lonard, D. M., Cowser, L. M., Bennett, C. F., O'Malley, B. W. and Smith, C. L. (2002) Reduction of coactivator expression by antisense oligodeoxynucleotides inhibits ERalpha transcriptional activity and MCF-7 proliferation. *Mol Endocrinol* **16**, 253-70. [PubMed](#) | [Back to Text](#)

Debes, J. D., Schmidt, L. J., Huang, H. and Tindall, D. J. (2002) p300 mediates androgen-independent transactivation of the androgen receptor by interleukin 6. *Cancer Res* **62**, 5632-6. [PubMed](#) | [Back to Text](#)

Elbashir, S. M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K. and Tuschl, T. (2001a) Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* **411**, 494-8. [PubMed](#) | [Back to Text](#)

Elbashir, S. M., Lendeckel, W. and Tuschl, T. (2001b) RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev* **15**, 188-200. [PubMed](#) | [Back to Text](#)

Elbashir, S. M., Martinez, J., Patkaniowska, A., Lendeckel, W. and Tuschl, T. (2001c) Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo-lysate. *Embo J* **20**, 6877-88. [PubMed](#) | [Back to Text](#)

Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E. and Mello, C. C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**, 806-11. [PubMed](#) | [Back to Text](#)

Hannon, G. J. (2002) RNA interference. *Nature* **418**, 244-51. [PubMed](#) | [Back to Text](#)

Li, X., Wong, J., Tsai, S. Y., Tsai, M. J. and O'Malley, B. W. (2003) Progesterone and glucocorticoid receptors recruit distinct coactivator complexes and promote distinct patterns of local chromatin modification. *Mol Cell Biol* **23**, 3763-73. [PubMed](#) | [Back to Text](#)

McManus, M. T., Petersen, C. P., Haines, B. B., Chen, J. and Sharp, P. A. (2002) Gene silencing using micro-RNA designed hairpins. *Rna* **8**, 842-50. [PubMed](#) | [Back to Text](#)

Paddison, P. J., Caudy, A. A., Bernstein, E., Hannon, G. J. and Conklin, D. S. (2002) Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes Dev* **16**, 948-58. [PubMed](#) | [Back to Text](#)

Plass, J. R., Mol, O., Heegsma, J., Geuken, M., Faber, K. N., Jansen, P. L. and Muller, M. (2002) Farnesoid X receptor and bile salts are involved in transcriptional regulation of the gene encoding the human bile salt export pump. *Hepatology* **35**, 589-96. [PubMed](#) | [Back to Text](#)

Rahman, M. M., Miyamoto, H., Lardy, H. and Chang, C. (2003a) Inactivation of androgen receptor coregulator ARA55 inhibits androgen receptor activity and agonist effect of antiandrogens in prostate cancer cells. *Proc Natl Acad Sci U S A* **100**, 5124-9. [PubMed](#) | [Back to Text](#)

Rahman, M. M., Miyamoto, H., Takatera, H., Yeh, S., Altuwaijri, S. and Chang, C. (2003b) Reducing the agonist activity of antiandrogens by a dominant-negative androgen receptor coregulator ARA70 in prostate cancer cells. *J Biol Chem* **278**, 19619-26. [PubMed](#) | [Back to Text](#)

Schreiber, S. N., Knutti, D., Brogli, K., Uhlmann, T. and Kralli, A. (2003) The transcriptional coactivator PGC-1 regulates the expression and activity of the orphan nuclear receptor estrogen-related receptor  $\alpha$  (ERR $\alpha$ ). *J Biol Chem* **278**, 9013-8. [PubMed](#) | [Back to Text](#)

Shang, Y. and Brown, M. (2002) Molecular determinants for the tissue specificity of SERMs. *Science* **295**, 2465-8. [PubMed](#) | [Back to Text](#)

Shinagawa, T. and Ishii, S. (2003) Generation of Ski-knockdown mice by expressing a long double-strand RNA from an RNA polymerase II promoter. *Genes*

*Dev* **17**, 1340-5. PubMed | Back to Text

Sui, G., Soohoo, C., Affar el, B., Gay, F., Shi, Y. and Forrester, W. C. (2002) A DNA vector-based RNAi technology to suppress gene expression in mammalian cells. *Proc Natl Acad Sci U S A* **99**, 5515-20. PubMed | Back to Text

Williams, B. R. (1997) Role of the double-stranded RNA-activated protein kinase (PKR) in cell regulation. *Biochem Soc Trans* **25**, 509-13. PubMed | Back to Text

Wright, M. E., Tsai, M. J. and Aebersold, R. (2003) Androgen receptor represses the neuroendocrine transdifferentiation process in prostate cancer cells. *Mol Endocrinol* **17**, 1726-37. PubMed | Back to Text

Yoon, H. G., Chan, D. W., Huang, Z. Q., Li, J., Fondell, J. D., Qin, J. and Wong, J. (2003) Purification and functional characterization of the human N-CoR complex: the roles of HDAC3, TBL1 and TBLR1. *Embo J* **22**, 1336-46. PubMed | Back to Text

Zamore, P. D., Tuschl, T., Sharp, P. A. and Bartel, D. P. (2000) RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* **101**, 25-33. PubMed | Back to Text

Zhou, G., Hashimoto, Y., Kwak, I., Tsai, S. Y. and Tsai, M. J. (2003) Role of the steroid receptor coactivator SRC-3 in cell growth. *Mol Cell Biol* **23**, 7742-55. PubMed | Back to Text

ISSN# 1550-7629

© 2003 - 2005 NURSA, The Nuclear Receptor Signaling Atlas.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**